## Associated Foot Surgeons Patient Surgical Instructions



Patient Name:							
	Instructions Before Surgery						
	If you are being treated for fluid retention, hypertension, diabetes, and/or a heart condition we will request:   Surgery Clearance Letter (from your primary, cardiologist, or other specialist(s).  Chest X-Ray  Stop NSAID's (Aspirin/Motrin/Advil/Aleve, ect 7 days prior to surgery)  Please contact your provider/cardiologist to find out their requirement for surgery clearance.  The tests/clearance documents must be submitted to our office, no later than 7 days prior to your ashedulad surgery.						
	scheduled surgery.  DO NOT EAT OR DRINK after am the day of surgery.  Surgery Center: A nurse from the surgery center will contact you a few days before surgery. They will review your medications, advise you on what medications to take the morning of surgery, update you on any changes and provide the arrival timeam/pm.						
	Hospital: Phone Pre Testing or Pre Testing Appointment: You must make transportation arrangements. If you are not familiar with your "Outpatient Surgery" benefits, we recommend you contact your insurance company to find out what you are responsible for paying.  Upon receipt of the insurance payment your portion (if any) must be paid within 30 days.  If you need crutches, we recommend practicing with them ahead of time. You can get them at any pharmacy without a prescription.  Hibiclens is a Pre-Surgical Scrub that can be found at Walgreens or Walmart. Please scrub your foot/feet from your toes to your knee the night before and the morning of the surgery.  Remove all nail polish. (Clear polish is acceptable) Wear clothing that will fit over a bulky bandage. Wear comfortable low heeled or no-heeled shoes.						
	POST OP INSTRUCTIONS  □ Please read all instructions prior to surgery						
	***If you have a fever and/or develop pain in the calf, go to your nearest emergency room.*** You have had a surgical operation. Sometimes limited amount of discomfort and swelling is expected. In some cases the skin may take on a bruised appearance. This is no cause for alarm. On the day of surgery, keep the foot elevated with an ice pack over the surgical site, except to go to the bathroom. Do not use hot water bags or electric pads. When walking or standing, always wear your surgical shoe/boot. A slight amount of blood may appear on the dressing. Elevate the foot above the level of the heart and apply an ice pack. If there is active or persistent bleeding, call our office at once.						



	If discomfort is experienced when the anesthesia wears off, take prescribed pain medication or an over the counter pain mediation of your choice. Do not take the medication more often than prescribed and do not drink alcoholic beverages while under the effect of the prescribed pain medication. If you are taking any narcotic pain medication, we recommend increasing your fluid intake and fiber. Keep the dressing completely dry. If it accidentally gets wet, notify our office. Do not remove the dressing at any time.  Beginning the day after surgery, move your foot up and down five minutes per hour to help prevent the formation of a blood clot in your calf muscle.  If any of the following occurs contact our office immediately: Redness with a white fluid discharge, an increase in body temperature, and/or nausea/vomiting.  Following the above directions will aid your recovery, however, if redness occurs and a white fluid is dis charged or a fever, please consult our office immediately.						
		S P	<b>urgery Schedul</b> o hone: 618-277-5	act Information: er: Kristen 700 x 5 Fax: 877- sociatedfoot.com	318-4562		
☐ Belleville Surgical Center  28 North 64th Street Belleville, IL 62223 Phone: 618-398-5705 www.bellevillesurgical.com Pre-Register Online			☐ Anderson Surgery Center 6800 Illinois Rt 162 Maryville, IL 62062 Phone: 618-288-5711 www.andersonhospital.org				
St Elizabeth's Hospital 211 South Third Street Belleville, IL 62220 Phone: 618-234-2120 www.steliz.org				☐ Memorial Hospital 4500 Memorial Drive Belleville, IL 62226 Phone: 618-233-7750 www.memhosp.com			
results	must be	You must contact you forwarded to our office cheduled on:	e no later than	7 day prior to you	r scheduled sur	gery date.	
Phy	sician:	Dr. Whittenburg	Dr. Snook	Dr. Reichert	Dr. Dugan	Dr. Hart	
	While in	☐ Cast/splint clean and your cast, we highly reaff you cannot take Aspi	dry. commend takin			ntil your cast is re	



Ш	Protect cast/splint with plastic bag during rain or snow.
	A wet cast/splint must be changed within 24 hours, to prevent skin breakdown.
	Keep the skin around the edge of a cast/splint clean and dry.
	Range of motion and elevation are important to control swelling.
	1. Bend and straighten the injured extremity several times daily.
	2. Elevate the injured extremity above your heart 20-30 minutes several times daily or whenever you have
	any of the following symptoms:
	An increase in swelling, pressure, or tightness inside the cast/splint
	An increase in pain tingling or numbness
	A decrease in circulation at the end of the extremity
	A cast is meant to be snug and to immobilize the involved joints; however, it should not cut off or
	decrease circulation of the extremity.
	DO NOT put anything inside of your cast/splint, or pull out any of the padding.
	Do not remove the cast/splint yourself.
	☐ Please report any of the following symptoms to your physician, within 24 hours:
	An increase in pain, pressure, swelling, tingling or numbness that cannot be controlled with
	elevation and range of motion.
	• An increase in drainage, foul odor, or if there is an increase in body temperature, nausea, or
	vomiting.
	<ul> <li>Any bandage or cast that gets wet, broken, cracked, too loose, or if cast/splint is causing pain</li> </ul>
	due to rubbing.
You	ur post op appointment is scheduled on: at